



STABILITY OF EULER-LAGRANGE-JENSEN'S (a, b) - SEXTIC FUNCTIONAL EQUATION IN MULTI-BANACH SPACES

JOHN MICHAEL RASSIAS¹, R. MURALI^{2,*} AND A. ANTONY RAJ²

¹*Pedagogical Department E.E., National and Kapodistrian University of Athens, Section of Mathematics and Informatics, Athens, 15342, Greece*

²*PG and Research Department of Mathematics, Sacred Heart College (Autonomous), Tirupattur - 635 601, Tamil Nadu, India*

**Corresponding author: shcrmurali@yahoo.co.in*

ABSTRACT. In this paper, we prove the Hyers-Ulam Stability of Euler-Lagrange-Jensen's (a, b) -Sextic Functional Equation in Multi-Banach Spaces.

1. INTRODUCTION AND PRELIMINARIES

The theory of stability is an important branch of the qualitative theory of functional equations. The concept of stability for a functional equation arises when one replaces a functional equation by an inequality which acts as a perturbation of the equation. The first stability problem of functional equation was raised by S.M. Ulam [17] about seventy seven years ago. Since then, this question has attracted the attention of many researchers. Note that the affirmative solution to this question was given in the next year by D.H. Hyers [5] in 1941. In the year 1950, T. Aoki [1] generalized Hyers theorem for additive mappings. The result of Hyers was generalized independently by Th.M.Rassias [14] for linear mappings by considering an unbounded Cauchy difference. In 1994, a further generalization of Th.M. Rassias theorem was obtained by P.Gavruta [4].

Then, the stability problem of several functional equations has been extensively investigated by a number

Received 2017-10-25; accepted 2018-01-04; published 2018-03-07.

2010 *Mathematics Subject Classification.* 39B52, 39B72, 39B82.

Key words and phrases. Hyers-Ulam stability; multi-Banach Spaces; sextic functional equation; direct method.

©2018 Authors retain the copyrights of their papers, and all open access articles are distributed under the terms of the Creative Commons Attribution License.

of authors, and there are many interesting results concerning this problem ([3, 6, 7, 9, 11–13, 15, 16, 18, 19]). The Hyers-Ulam stability of functional equation is investigated and the investigation is following. Here, we establish the Hyers-Ulam Stability of Euler-Lagrange-Jensen’s (a, b) - Sextic Functional Equation is of the form

$$\begin{aligned}
 & f(ax + by) + f(bx + ay) + (a - b)^6 \left[f\left(\frac{ax - by}{a - b}\right) + f\left(\frac{bx - ay}{b - a}\right) \right] \\
 & = 64(ab)^2(a^2 + b^2) \left[f\left(\frac{x + y}{2}\right) + f\left(\frac{x - y}{2}\right) \right] + 2(a^2 - b^2)(a^4 - b^4) [f(x) + f(y)]
 \end{aligned} \tag{1.1}$$

where $a \neq b$ such that $k \in \mathbb{R}$, $h = a + b \neq 0, \pm 1$ in Multi-Banach Spaces by using direct and fixed point method.

Definition 1.1. [2] A Multi- norm on $\{\mathcal{A}^k : k \in \mathbb{N}\}$ is a sequence $(\|\cdot\|) = (\|\cdot\|_k : k \in \mathbb{N})$ such that $\|\cdot\|_k$ is a norm on \mathcal{A}^k for each $k \in \mathbb{N}$, $\|x\|_1 = \|x\|$ for each $x \in \mathcal{A}$, and the following axioms are satisfied for each $k \in \mathbb{N}$ with $k \geq 2$:

- (1) $\|(x_{\sigma(1)}, \dots, x_{\sigma(k)})\|_k = \|(x_1 \dots x_k)\|_k$, for $\sigma \in \Psi_k, x_1, \dots, x_k \in \mathcal{A}$;
- (2) $\|(\alpha_1 x_1, \dots, \alpha_k x_k)\|_k \leq (\max_{i \in \mathbb{N}_k} |\alpha_i|) \|(x_1 \dots x_k)\|_k$
for $\alpha_1 \dots \alpha_k \in \mathbb{C}, x_1, \dots, x_k \in \mathcal{A}$;
- (3) $\|(x_1, \dots, x_{k-1}, 0)\|_k = \|(x_1, \dots, x_{k-1})\|_{k-1}$, for $x_1, \dots, x_{k-1} \in \mathcal{A}$;
- (4) $\|(x_1, \dots, x_{k-1}, x_{k-1})\|_k = \|(x_1, \dots, x_{k-1})\|_{k-1}$ for $x_1, \dots, x_{k-1} \in \mathcal{A}$.

In this case, we say that $(\mathcal{A}^k, \|\cdot\|_k) : k \in \mathbb{N}$ is a multi - normed space.

Suppose that $(\mathcal{A}^k, \|\cdot\|_k) : k \in \mathbb{N}$ is a multi - normed space, and take $k \in \mathbb{N}$. We need the following two properties of multi - norms. They can be found in [2].

- (a) $\|(x, \dots, x)\|_k = \|x\|$, for $x \in \mathcal{A}$,
- (b) $\max_{i \in \mathbb{N}_k} \|x_i\| \leq \|(x_1, \dots, x_k)\|_k \leq \sum_{i=1}^k \|x_i\| \leq k \max_{i \in \mathbb{N}_k} \|x_i\|, \forall x_1, \dots, x_k \in \mathcal{A}$.

It follows from (b) that if $(\mathcal{A}, \|\cdot\|)$ is a Banach space, then $(\mathcal{A}^k, \|\cdot\|_k)$ is a Banach space for each $k \in \mathbb{N}$. In this case, $(\mathcal{A}^k, \|\cdot\|_k) : k \in \mathbb{N}$ is a multi - Banach space.

2. STABILITY OF FUNCTIONAL EQUATION (1.1) IN MULTI-BANACH SPACES: DIRECT METHOD

Theorem 2.1. Let X be a linear space and $(Y^n, \|\cdot\|_n) : n \in \mathbb{N}$ be a multi-Banach Spaces. Let $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ such that

$$\sup_{k \in \mathbb{N}} \|(\mathcal{D}f(x_1, y_1), \dots, \mathcal{D}f(x_k, y_k))\|_k \leq \epsilon \tag{2.1}$$

$\forall x_1, \dots, x_k, y_1, \dots, y_k \in Y$. Then there exists a unique sextic mapping $S : X \rightarrow Y$ such that

$$\sup_{k \in \mathbb{N}} \|(f(x_1) - S(x_1), \dots, f(x_k) - S(x_k))\| \leq \frac{\epsilon}{h^6} \tag{2.2}$$

Proof. Letting $y_i = x_i$ where $i = 1, 2, \dots, k$ in (2.1), we arrive at

$$\sup_{k \in \mathbb{N}} \left\| \left(\frac{1}{h^6} f(hx_1) - f(x_1), \dots, \frac{1}{h^6} f(hx_k) - f(x_k) \right) \right\| \leq \frac{\epsilon}{2h^6} \tag{2.3}$$

Now, Replacing x_i by $2x_i$ where $i = 1, 2, \dots, k$ and dividing by 2 in above equation, we get

$$\sup_{k \in \mathbb{N}} \left\| \left(\frac{f(2hx_1)}{h^6} - f(x_1), \dots, \frac{f(2hx_k)}{h^6} - f(x_k) \right) \right\| \leq \frac{\epsilon}{2^2 h^6} + \frac{\epsilon}{2h^6} \tag{2.4}$$

By using induction for a positive integer n , we obtain

$$\sup_{k \in \mathbb{N}} \left\| \left(\frac{f(2^n hx_1)}{2^n h^6} - f(x_1), \dots, \frac{f(2^n hx_k)}{2^n h^6} - f(x_k) \right) \right\| \leq \frac{1}{h^6} \sum_{i=0}^{n-1} \frac{\epsilon}{2^{i+1}} \leq \frac{1}{h^6} \sum_{i=0}^{\infty} \frac{\epsilon}{2^{i+1}} \tag{2.5}$$

Now, we have to show that the sequence $\left\{ \frac{f(2^n hx)}{2^n h^6} \right\}$ is a Cauchy sequence, by fixing $x \in X$ and replacing x_1, \dots, x_k by $x, 2x, \dots, 2^{k-1}x$ such that

$$\begin{aligned} & \sup_{k \in \mathbb{N}} \left\| \left(\frac{f(2^n hx)}{2^n h^6} - \frac{f(2^m x)}{2^m}, \dots, \frac{f(2^{n+k-1} hx)}{2^{n+k-1} h^6} - \frac{f(2^{m+k-1} x)}{2^{m+k-1}} \right) \right\| \\ & \leq \sup_{k \in \mathbb{N}} \left\| \left(\frac{f(2^n hx)}{2^n h^6} - \frac{f(2^m x)}{2^m}, \dots, \frac{1}{2^{k-1}} \left[\frac{f(2^n (2^{k-1} hx))}{2^n h^6} - \frac{f(2^m (2^{k-1} x))}{2^m} \right] \right) \right\| \end{aligned}$$

Using the definition of Multi-norm, we arrive at

$$\begin{aligned} & \sup_{k \in \mathbb{N}} \left\| \left(\frac{f(2^n hx)}{2^n h^6} - \frac{f(2^m x)}{2^m}, \dots, \frac{f(2^n (2^{k-1} hx))}{2^n h^6} - \frac{f(2^m (2^{k-1} x))}{2^m} \right) \right\| \\ & \leq \frac{1}{h^6} \sum_{i=m}^{n-1} \frac{\epsilon}{2^{i+1}}. \end{aligned} \tag{2.6}$$

Hence the above inequality (2.6), shows that $\left\{ \frac{f(2^n hx)}{2^n h^6} \right\}$ is a Cauchy sequence as $n \rightarrow \infty$. Since Y is complete, then the sequence $\left\{ \frac{f(2^n hx)}{2^n h^6} \right\}$ converges to a fixed point $S(x) \in Y$ such that

$$S(x) = \lim_{n \rightarrow \infty} \frac{f(2^n hx)}{2^n h^6}.$$

Therefore, as $n \rightarrow \infty$, the inequality (2.5) implies the inequality (2.2). Obviously, one can find the uniqueness of the mapping $S : X \rightarrow Y$, using the definition of multi-norm. That is, we can prove $S = S'$. \square

Corollary 2.1. Let X be a linear space and $((Y^n, \|\cdot\|_n) : n \in \mathbb{N})$ be a multi-Banach space. Let $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ such that

$$\sup_{k \in \mathbb{N}} \|Df(x_1, y_1, \dots, x_k, y_k)\|_k \leq \phi(x_1, y_1, \dots, x_k, y_k) \tag{2.7}$$

for all $x_1, \dots, x_k, y_1, \dots, y_k \in X$. Then there exists a unique sextic mapping $S : X \rightarrow Y$ such that

$$\sup_{k \in \mathbb{N}} \|f(x_1) - S(x_1), \dots, f(x_k) - S(x_k)\|_k \leq \frac{1}{h^6} \sum_{i=1}^{\infty} \frac{1}{2^{i+1}} \phi(2^i x_1, x_1, \dots, 2^i x_k, x_k) \tag{2.8}$$

for all $x_1, \dots, x_k \in X$.

Proof. Proof is similar to that of Theorem 2.1 by replacing the condition $\phi(x_1, y_1, \dots, x_k, y_k)$ in place of ϵ . \square

Corollary 2.2. Let X be a linear space and $((Y^n, \|\cdot\|_n) : n \in \mathbb{N})$ be a multi-Banach space. Let $0 < p < 6$, $\theta \geq 0$ and $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ such that

$$\sup_{k \in \mathbb{N}} \|Df(x_1, y_1, \dots, x_k, y_k)\|_k \leq \theta (\|x_1\|^p + \|y_1\|^p, \dots, \|x_k\|^p + \|y_k\|^p) \tag{2.9}$$

for all $x_1, \dots, x_k, y_1, \dots, y_k \in X$. Then there exists a unique sextic mapping $S : X \rightarrow Y$ such that

$$\sup_{k \in \mathbb{N}} \|f(x_1) - S(x_1), \dots, f(x_k) - S(x_k)\|_k \leq \frac{\theta}{h^6(2^p - 1)} (\|x_1\|^p, \dots, \|x_k\|^p) \tag{2.10}$$

for all $x_1, \dots, x_k \in X$.

Proof. Proof is similar to that of Theorem 2.1 by replacing the condition

$\theta (\|x_1\|^p + \|y_1\|^p, \dots, \|x_k\|^p + \|y_k\|^p)$ in place of ϵ . \square

3. STABILITY OF FUNCTIONAL EQUATION (1.1) IN MULTI-BANACH SPACES: FIXED POINT METHOD

Theorem 3.1. Let X be a linear space and $((Y^n, \|\cdot\|_n) : n \in \mathbb{N})$ be a multi-Banach Spaces. Let $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ such that

$$\sup_{k \in \mathbb{N}} \|(\mathcal{D}f(x_1, y_1), \dots, \mathcal{D}f(x_k, y_k))\|_k \leq \epsilon \tag{3.1}$$

$\forall x_1, \dots, x_k, y_1, \dots, y_k \in Y$. Then there exists a unique sextic mapping $S : X \rightarrow Y$ such that

$$\sup_{k \in \mathbb{N}} \|(f(x_1) - S(x_1), \dots, f(x_k) - S(x_k))\| \leq \frac{\epsilon}{2(h^6 - 1)} \tag{3.2}$$

Proof. Letting $y_i = x_i$ where $i = 1, 2, \dots, k$ in (2.1), we arrive at

$$\sup_{k \in \mathbb{N}} \left\| \left(\frac{1}{h^6} f(hx_1) - f(x_1), \dots, \frac{1}{h^6} f(hx_k) - f(x_k) \right) \right\| \leq \frac{\epsilon}{2h^6} \tag{3.3}$$

Let $\Psi = \{l : X \rightarrow Y | l(0) = 0\}$ and introduce the generalized metric d defined on Ψ by

$$d(l, m) = \inf \left\{ \Psi \in [0, \infty] \mid \sup_{k \in \mathbb{N}} \|l(x_1) - m(x_1), \dots, l(x_k) - m(x_k)\|_k \leq \Psi \quad \forall x_1, \dots, x_k \in X \right\}$$

Then it is easy to show that Ψ, d is a generalized complete metric space, See [8].

We define an operator $\mathcal{J} : \Psi \rightarrow \Psi$ by

$$\mathcal{J}l(x) = \frac{1}{h^6} l(hx) \quad x \in X$$

We assert that \mathcal{J} is a strictly contractive operator. Given $l, m \in \Psi$, let $\Psi \in [0, \infty]$ be an arbitrary constant with $d(l, m) \leq \Psi$. From the definition it follows that

$$\sup_{k \in \mathbb{N}} \|l(x_1) - m(x_1), \dots, l(x_k) - m(x_k)\|_k \leq \Psi \quad x_1, \dots, x_k \in X.$$

Therefore, $\sup_{k \in \mathbb{N}} \|(\mathcal{J}l(x_1) - \mathcal{J}m(x_1), \dots, \mathcal{J}l(x_k) - \mathcal{J}m(x_k))\|_k \leq \frac{1}{h^6} \Psi$
 $x_1, \dots, x_k \in X$. Hence, it holds that

$$d(\mathcal{J}l, \mathcal{J}m) \leq \frac{1}{h^6} \Psi d(\mathcal{J}l, \mathcal{J}m) \leq \frac{1}{h^6} d(l, m)$$

$\forall l, m \in \Psi$.

This means that \mathcal{J} is strictly contractive operator on Ψ with the Lipschitz constant $L = \frac{1}{h^6}$.

By (3.3), we have $d(\mathcal{J}f, f) \leq \frac{\epsilon}{2h^6}$. Applying the Theorem 2.2 in [10], we deduce the existence of a fixed point of \mathcal{J} that is the existence of mapping $S : X \rightarrow Y$ such that

$$S(hx) = h^6 S(x) \quad \forall x \in X.$$

Moreover, we have $d(\mathcal{J}^n f, S) \rightarrow 0$, which implies

$$S(x) = \lim_{n \rightarrow \infty} \mathcal{J}^n f(x) = \lim_{n \rightarrow \infty} \frac{f(h^n x)}{h^{6n}}$$

for all $x \in X$.

Also, $d(f, S) \leq \frac{1}{1 - \mathcal{L}} d(\mathcal{J}f, f)$ implies the inequality

$$\leq \frac{\epsilon}{2(h^6 - 1)}.$$

Doing $x_1 = \dots = x_k = h^n x$, and $y_1 = \dots = y_k = h^n y$ in (1.1) and dividing by h^{6n} . Now, applying the property (a) of multi-norms, we have

$$\begin{aligned} \|DS(x, y)\| &= \lim_{n \rightarrow \infty} \frac{1}{h^{6n}} \|Df(h^n x, h^n y)\| \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{h^{6n}} = 0 \end{aligned}$$

for all $x, y \in X$. The uniqueness of S follows from the fact that S is the unique fixed point of \mathcal{J} with the property that there exists $\ell \in (0, \infty)$ such that

$$\sup_{k \in \mathbb{N}} \|(f(x_1) - S(x_1), \dots, f(x_k) - S(x_k))\|_k \leq \ell$$

for all $x_1, \dots, x_k \in X$.

Hence the proof. □

Corollary 3.1. Let X be a linear space and $((Y^n, \|\cdot\|_n) : n \in \mathbb{N})$ be a multi-Banach space. Let $0 < p < 6$, $\theta \geq 0$ and $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ such that

$$\sup_{k \in \mathbb{N}} \|Df(x_1, y_1, \dots, x_k, y_k)\|_k \leq \theta (\|x_1\|^p + \|y_1\|^p, \dots, \|x_k\|^p + \|y_k\|^p) \quad (3.4)$$

for all $x_1, \dots, x_k, y_1, \dots, y_k \in X$. Then there exists a unique sextic mapping $S : X \rightarrow Y$ such that

$$\sup_{k \in \mathbb{N}} \|f(x_1) - S(x_1), \dots, f(x_k) - S(x_k)\|_k \leq \frac{2\theta}{h^6 - 2h^p} (\|x_1\|^p, \dots, \|x_k\|^p) \quad (3.5)$$

for all $x_1, \dots, x_k \in X$.

Proof. Proof is similar to that of Theorem 3.1 by replacing the condition

$\theta (\|x_1\|^p + \|y_1\|^p, \dots, \|x_k\|^p + \|y_k\|^p)$ in place of ϵ . □

Corollary 3.2. Let X be a linear space and $((Y^n, \|\cdot\|_n) : n \in \mathbb{N})$ be a multi-Banach space. Let $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ such that

$$\sup_{k \in \mathbb{N}} \|Df(x_1, y_1, \dots, x_k, y_k)\|_k \leq \phi(x_1, y_1, \dots, x_k, y_k) \quad (3.6)$$

for all $x_1, \dots, x_k, y_1, \dots, y_k \in X$. Then there exists a unique sextic mapping $S : X \rightarrow Y$ such that

$$\sup_{k \in \mathbb{N}} \|f(x_1) - S(x_1), \dots, f(x_k) - S(x_k)\|_k \leq \frac{1}{2(h^6 - 1)} \phi(x_1, x_1, \dots, x_k, x_k) \quad (3.7)$$

for all $x_1, \dots, x_k \in X$.

Proof. Proof is similar to that of Theorem 3.1 by replacing the condition $\phi(x_1, y_1, \dots, x_k, y_k)$ in place of ϵ . □

REFERENCES

- [1] T. Aoki, On the stability of the linear transformation in Banach spaces, J.Math. Soc. Japan 2 (1950), 64-66.
- [2] Dales, H.G and Moslehian, Stability of mappings on multi-normed spaces, Glasgow Math. J. 49 (2007), 321-332.
- [3] Fridoun Moradlou, Approximate Euler-Lagrange-Jensen type Additive mapping in Multi-Banach Spaces: A Fixed point Approach, Commun. Korean Math. Soc. 28 (2013), 319-333.
- [4] P. Gavruta, On a problem of G.Isac and Th.M.Rassias concerning the stability of mappings, J. Math. Anal. Appl. 261 (2001), 543-553.
- [5] D.H.Hyers, On the stability of the linear functional equation, Proc. Nat. Acad. Sci. USA 27 (1941), 222-224.
- [6] John Michael Rassias, R. Murali, Matina John Rassias, A. Antony Raj, General Solution, stability and Non-stability of Quattuorvigintic functional equation in Multi-Banach spaces, Int. J. Math. Appl. 5 (2017), 181-194.
- [7] Manoj Kumar and Ashish Kumar, Stability of Jensen Type Quadratic Functional Equations in Multi-Banach Spaces, Int. J. Math. Arch. 3(4) (2012), 1372-1378.
- [8] D. Mihet and V. Radu, On the stability of the additive Cauchy functional equation in random normed spaces, J. Math. Anal. Appl. 343 (2008), 567-572.
- [9] R. Murali, Matina J. Rassias and V. Vithya, The General Solution and stability of Nonadecic Functional Equation in Matrix Normed Spaces, Malaya J. Mat. 5(2) (2017), 416-427.
- [10] V.Radu, The fixed point alternative and the stability of functional equations, Fixed Point Theory 4 (2003), 91-96.

-
- [11] K. Ravi, J.M. Rassias and B.V. Senthil Kumar, Ulam-Hyers stability of undecic functional equation in quasi-*beta* normed spaces fixed point method, Tbilisi Math. Sci. 9 (2) (2016), 83-103.
- [12] K. Ravi, J.M. Rassias, S. Pinelas and S.Suresh, General solution and stability of Quattuordecic functional equation in quasi *beta* normed spaces, Adv. Pure Math. 6 (2016), 921-941.
- [13] J. M. Rassias and M. Eslamian, Fixed points and stability of nonic functional equation in quasi- β -normed spaces, Cont. Anal. Appl. Math. 3 (2) (2015), 293-309.
- [14] Th.M. Rassias, On the stability of the linear mappings in Banach Spaces, Proc. Amer. Math. Soc., 72 (1978), 297-300.
- [15] Th.M. Rassias, On the stability of functional equations and a problem of Ulam, Acta. Appl. Math. 62 (2000) 23 - 130.
- [16] Tian Zhou Xu, John Michael Rassias and Wan Xin Xu, Generalized Ulam - Hyers Stability of a General Mixed AQCQ functional equation in Multi-Banach Spaces: A Fixed point Approach, Eur. J. Pure Appl. Math. 3 (2010), 1032-1047.
- [17] S.M. Ulam, A Collection of the Mathematical Problems, Interscience, New York, (1960).
- [18] T. Z. Xu, J. M. Rassias, M. J. Rassias and W. X. Xu, A fixed point approach to the stability of quintic and sextic functional equations in quasi- β -normed spaces, J. Inequal. Appl. 2010 (2010), Article ID 423231.
- [19] Xiuzhong Wang, Lidan Chang, Guofen Liu, Orthogonal Stability of Mixed Additive-Quadratic Jensen Type Functional Equation in Multi-Banach Spaces, Adv. Pure Math. 5 (2015), 325-332.